

CLENPIQ™
 (sodium picosulfate, magnesium oxide,
 and anhydrous citric acid) Oral Solution
 10 mg/3.5 g/12 g per 160 mL bottle

TAKING CLENPIQ

STEPS TO COMPLETE THE PREP

Day-Before-Dose Regimen

Afternoon/Early Evening Before
 BETWEEN 4-6 PM

4pm

Drink 40 oz (5 cups) of clear liquids.*
 Finish liquids over the next 5 hours.

Evening Before
 Between 10 PM - 12 AM

9pm

Drink at least 24 oz (3 cups) of clear liquids over the next 5 hours.
 Finish liquids 2 hours before your colonoscopy
 or as advised by your doctor.



Hydration is important and it's part of the prep. Make sure to hydrate before you take the prep, while you're taking the prep, and after the prep.

Prep Assistant Use the prep assistant as a guide for completing your prep.

The date and time of your colonoscopy is _____

You can keep hydrating with clear liquids until Midnight Before procedure.

ON THE DAY BEFORE THE COLONOSCOPY, stop eating all solid food and dairy, and start hydrating by drinking clear liquids.



Afternoon/Early Evening before your colonoscopy

4-6 PM

Drink one bottle of CLENPIQ
 Drink CLENPIQ right from the bottle

Then, keep hydrating
 Drink 5 cups (8 oz each) of clear liquid.
 Check off the cups below as you go!



Start: _____
 Finish: _____

Evening before your colonoscopy

10 PM - 12 AM

Drink the other bottle of CLENPIQ
 Drink CLENPIQ right from the bottle

Then, keep hydrating
 Drink at least 3 cups (8 oz each) of clear liquid.
 Check off the cups below as you go!



Start: _____
 Finish: _____

*After your first dose, if severe bloating, swelling, or stomach pain occurs, delay the second dose until the symptoms resolve.
 Please see reverse page for additional information about CLENPIQ, the importance of a clear liquid diet, and hydration.

INDICATION

CLENPIQ is a prescription medicine used by adults to clean the colon before a colonoscopy. CLENPIQ cleans your colon by causing you to have diarrhea. Cleaning your colon helps your healthcare provider see the inside of your colon more clearly during your colonoscopy.

IMPORTANT SAFETY INFORMATION

- Do not take CLENPIQ if your healthcare provider has told you that you have serious kidney problems, a blockage in your intestine (bowel obstruction), an opening in the wall of your stomach or intestines (bowel perforation), a very dilated intestine (toxic megacolon), problems with the emptying of food and fluid from your stomach (gastric retention), or an allergy to any of the ingredients in CLENPIQ.
- CLENPIQ and other bowel preparations can cause serious side effects, including serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause abnormal heartbeats that may result in death, seizures (this can happen even if you have never had a seizure), or kidney problems. Your chance of having fluid loss and changes in blood salts with CLENPIQ is higher if you have heart problems, have kidney problems, or take water pills or non-steroidal anti-inflammatory drugs (NSAIDs).

See complete Important Safety Information on the back.

1 – Gastroenterology Specialists of Gwinnett PC

**721 Wellness Way NW, Suite 100, Lawrenceville, GA 30045 (770) 995-7989, (770) 995-3113
3855 Pleasant Hill Road, Suite 230, Duluth, GA 30096 (770) 495-8300**

**Zack Z. Martin, M.D. D. Bradley Lord, M.D. W. Gordon Tanner, Jr., M.D.
Jae W. Nam, M.D. Pamela Boyd, APRN FNP-C**

LOW FIBER/LOW RESIDUE AND LIQUID DIETS

LOW FIBER DIET: (3 DAYS PRIOR TO COLONOSCOPY):

Soups: Broth, cream soups, tomato, chicken and white rice, noodles

Meats/Protein: Any soft or easy to chew meat or fish. Such as fish or chicken. Avoid red meat.

Vegetables: Well-cooked or canned vegetables, without seeds or skins. Such as carrots, potatoes without skins and tomatoes without seeds or skins.

Fruits: All juices without pulp, cooked or canned fruits without seeds, membranes, or hulls including: applesauce, cherries, peaches, bananas, citrus fruits.

Cereals: No multigrain or high fiber cereals. Refined cereals or ready to eat oatmeal.

Starches: No whole grain, multigrain or wheat breads or pastas. May have enriched white bread, without seeds, biscuits, rolls, crackers or bread made from white flour, plain doughnuts or English muffins, plain macaroni, noodles or spaghetti, white rice, plain waffles or pancakes, saltines.

Cheese/Eggs: Cottage cheese, cream cheese, mild cheese such as American or Swiss. Yogurt without fruit. All eggs.

Desserts: Plain cake, cookies, gelatin, ice cream, pudding, sherbet, pie made from refined flour. No seeds, raisins, coconuts or nuts.

Beverages: Limit milk to two cups a day, carbonated beverages, coffee, tea, fruit-flavored drinks.

Fat: Crisp bacon, butter, cream, margarine, mayonnaise, oils, shortening, smooth salad dressing, sour cream.

LIQUID DIET: (All day on the day prior to the colonoscopy. NO SOLID FOOD FOR THE ENTIRE DAY.)

Soups: Clear consommé, bouillon or broth.

Beverages: Tea, coffee, carbonated drinks, water, Kool-Aid (no red or purple).

Juices: Cranberry (no red or purple), apple, strained lemonade, or orange juice.

Desserts: Gelatins, water-ices, popsicles (no red or purple).

Miscellaneous: Hard candy (Lifesavers, etc.)

1 – Gastroenterology Specialists of Gwinnett PC

**721 Wellness Way NW, Suite 100, Lawrenceville, GA 30045 (770) 995-7989, (770) 995-3113
3855 Pleasant Hill Road, Suite 230, Duluth, GA 30096 (770) 495-8300**

*Zack Z. Martin, M.D. D. Bradley Lord, M.D. W. Gordon Tanner, Jr., M.D.
Jae W. Nam, M.D. Pamela Boyd, APRN FNP-C*

COLONOSCOPY

Colonoscopy is an examination for the inside of the large intestine by means of a flexible tube attached to a source of bright light. The doctor is able to see the lining of the large intestine clearly through this tube.

After completing the preparation at home, come to the facility as instructed, where you will change into a gown. The test will be more fully explained, and you will be given an opportunity to ask questions. The nurse will then insert a small needle into your vein to administer the IV solution. You will receive medication through your IV before and during the procedure to make you relaxed and comfortable. During the exam, you will be lying on your left side on a stretcher bed.

After you are adequately sedated, the doctor will examine your rectum and insert the lubricated tube. During the procedure, you may experience crampy sensations similar to gas pains. These are due to the air put into your intestine through the tube in order to see well and to the tube passing around the bends of your intestine. You may feel that you need to move your bowels due to the presence of the tube and air. You are free to expel air as needed for your comfort. Any excess air can be suctioned out by the physician.

When the exam is over, you will rest in the recovery area until fully awake, and the doctor will explain the results of the exam to you and your family before you leave.

It is necessary to BRING A DRIVER FOR THE TRIP HOME, and you should not resume your usual work or driving until the day following the procedure. Do not eat or drink anything on the morning of the test unless specifically instructed otherwise. Please plan on taking your medications following the examination.

If you have any questions, please call our office at 770-995-3113 or 770-495-8300 even after hours.